

On Parrondo's paradox: how to construct unfair games by composing fair games

BY ERIC S. KEY¹, MALGORZATA M. KŁOSEK¹, DEREK ABBOTT²

¹*Department of Mathematical Sciences, University of Wisconsin Milwaukee,
Milwaukee WI 53201 USA*

²*Centre for Biomedical Engineering (CBME), Department of Electrical &
Electronic Engineering, University of Adelaide, Adelaide SA 5005 Australia*

We construct games of chance from simpler games of chance. We show that it may happen that the simpler games of chance are fair or unfavourable to a player and yet the new combined game is favourable – this is a counter-intuitive phenomenon known as Parrondo's paradox. We observe that all of the games in question are random walks in periodic environments (RWPE) when viewed on the proper time scale. Consequently, we use RWPE techniques to derive conditions under which Parrondo's paradox occurs.

Keywords: random walk in a periodic environment, random transport,
random games, Parrondo's paradox

1. Introduction

Parrondian strategies are where losing games can cooperate to win (Harmer & Abbott 1999a). The original example of Parrondo's games consist of two coin tossing games. Game A consists of Coin 1 biased to lose. Game B consists of two coins – Coin 2 with losing bias and Coin 3 with winning bias – but a state-dependent rule is chosen to favour the losing Coin 2. Hence both games A and B are losing games. However when A and B are alternated in a deterministic or even random manner, the player surprisingly has a winning expectation. This effect has been interpreted in terms of a discrete-time Brownian ratchet, at length, elsewhere (Harmer & Abbott 1999b) – where conventional Brownian ratchets (Doering 1995) have been the inspiration. An alternative view, we call the Boston interpretation (Stanley 1999, group discussion), recognises that although game B favours Coin 2 with losing bias, if the state-dependence is removed game B now favours the winning Coin 3 – then when games A and B are mixed, game A has the affect of randomisation or 'break-up' of game B's state-dependence, thus tilting favour towards Coin 3 with winning bias. This explanation was also independently deduced by J. Maynard Smith (1999, personal communication). Further to the ratchet interpretation and Boston interpretation, this paper will examine another viewpoint by considering the process as a random walk in a periodic environment (RWPE).

We now briefly summarise the literature on Parrondo's games. In (Harmer *et al.* 2000a) the state-dependent rule, for game B, is to choose Coin 2 if the player's capital is a multiple of some integer M – analysis showed the paradox could hold for general values of M . In (Pearce, 2000a) it is shown that the paradox can hold

when both games A and B are multiple coin games. In (Pearce 2000b) we have the first analysis of Parrondo's games in terms of Shannon entropy and in (Harmer *et al.* 2000b) the entropy parameter spaces are graphically displayed. The probability parameter space is shown in (Harmer *et al.* 2000c). In (Lee *et al.* 2000) a minimal three-state game with asymmetric transition probabilities is analysed and in (Parrondo *et al.* 2000) state-dependence on capital is replaced by dependence on the past history of the game, leading to a larger probability parameter space.

The surge of interest in analysing Parrondian games is motivated by a number of areas. Information theorists have long studied the problem of producing a fair game from biased coins (Gargamo and Vaccaro 1999) and the roots of this can be traced back to the work of von Neumann (1951) – Parrondo's games go a step further in producing a winning game from losing games. Seigman (1999, personal communication) has reinterpreted 'capital' of the games in terms electron occupancies of energy levels – the paradox can then be reproduced using the rate equation approach typically used in laser analysis. In the physical world there are many types of processes where losing helps to win, such as a sacrifice in the game of chess or a valley in the fitness landscape of an animal species. Many biological effects are linked to ratchet type phenomena and Westerhoff *et al.* (1986) have analysed enzyme transport with a four-state model. Applicability to population genetics, evolution and economics has been suggested (McClintock 1999). In finance, Maslov and Zhang (1998) have shown that under certain conditions capital can grow by investing in an asset with *negative* typical growth rate. Quantum ratchets have now been experimentally realised (Linke *et al.* 1999) and recasting Parrondian games, based on ratchet phenomena, as quantum games (Eisert *et al.* 1999; Goldenberg *et al.* 1999; Meyer 1999) is thus of interest.

In control theory, it can be shown that the combination of two unstable systems can become stable (Allison & Abbott 2000). Velocity of propagation through an array of coupled oscillators, under certain conditions, can *increase* even though the damping coefficient is *increased* (Sarmiento *et al.* 1999). In the area of granular flow, drift can occur in a counter-intuitive direction such as exemplified in the famous Brazil nut paradox (Rosato *et al.* 1987). Also declining branching processes can be combined to increase (Key 1987). Plaskota (1996) shows that noisy information can sometimes be better than clean information. In (Pinsky and Scheutzow 1992) it is shown that with switched diffusion processes in random media it is possible to get a positive-recurrent processes (*i.e.* with no drift) from mixed transient processes (*i.e.* with drifts all in the same direction) – this is almost certainly a continuous time analogue to the Parrondian discrete-time process. Assuming we construct Parrondo's games to only deal in transactions of one unit of capital per event, then we have a skip-free process, and a statistical interpretation of the central result is that declining birth-death processes can be combined to form an increase.

In this paper we further investigate Parrondo's paradox. We construct a class of composite games and investigate their fairness by formulating the problem in the language of random walks in periodic environments (RWPE). We find many new, interesting, and counter-intuitive results.

2. Mathematical tools

We construct a composite game from two simple games A and B . These two games can be combined in two ways: deterministically or stochastically.

If we have played n times, and n is divisible by k (for an integer k), we then play game A . If we have played n times and n is not divisible by k , we play game B . Thus games A and B are alternated in a deterministic pattern. We denote by Y_n our capital after n plays of this game.

To alternate games A and B randomly we toss a coin with probability p of heads. If the coin comes up heads, we play game A , and if tails, we play game B . We denote by Z_n be our capital after n repetitions.

The sequence of values of our capital in either of these games is a Markov random walk which changes by ± 1 in each epoch. The two random walks Z_n and Y_n differ in that Z_n is time homogeneous, while Y_n is not time homogeneous. Moreover, Z_n is a random walk in a periodic environment. The process Y_n is not a RWPE, but the process $Y'_n \equiv Y_{kn}$ is a RWPE.

As shown in §4 we extend this construction to define composite games from more than two simple games. In each case we identify the capital of the player as a RWPE.

We say that a game is fair, winning or losing if the random walk for the capital of a player, X_n , is recurrent or transient to ∞ , or to $-\infty$, respectively. That is, a Markov chain X_n is

$$\begin{aligned} &\text{recurrent (fair) if } P\{-\infty = \liminf_{n \rightarrow \infty} X_n < \limsup_{n \rightarrow \infty} X_n = \infty\} = 1; \\ &\text{transient to } \infty \text{ (winning) if } P\{\lim_{n \rightarrow \infty} X_n = +\infty\} = 1; \\ &\text{transient to } -\infty \text{ (losing) if } P\{\lim_{n \rightarrow \infty} X_n = -\infty\} = 1. \end{aligned}$$

We note that the characterisation of a game as fair, winning, or losing by the traditional comparisons $E[X_{n+1}|X_n] = X_n$, $E[X_{n+1}|X_n] > X_n$, and $E[X_{n+1}|X_n] < X_n$, respectively, does not cover the behaviour all random walks, in particular RWPE's.

(a) Key's criterion

We consider a time homogeneous random walk, X_n , in an N -periodic environment, or equivalently we have a state dependent random game $\mathbf{W}$. We assume that the maximal step size in the positive direction is R , while in the negative direction it is L , that is $P\{X_{n+1} \in \{-L + k, \dots, k + R\} | X_n = k\} = 1$. Moreover, for each k the maximum right and left step sizes are always possible, that is, $P\{X_{n+1} = -L + k | X_n = k\}P\{X_{n+1} = R + k | X_n = k\} > 0$. Given the environment, the walk X_n obeys the backward master equation

$$P\{\cdot | X_n = k\} = \sum_{j=-L}^R e(k, j)P\{\cdot | X_{n+1} = j + k\}, \quad (2.1)$$

where $e(k, j) \equiv P\{X_{n+1} = j + k | X_n = k\}$ denotes the transition probability from the state k to $j + k$ in one time epoch. We denote by $f_{k-i} \equiv P\{\cdot | X_{n+1} = k - i\}$

(since X_n is time homogeneous f does not depend on n) and rewrite equation (2.1) as a system for the vector $[f_{-L+k}, f_{-L+1+k}, \dots, f_{R-1+k}]^T$ with the matrix $\mathbf{A}_k$ whose entries given by

$$\mathbf{A}_k[i, j] = \begin{cases} -e(k, -L + j)/e(k, -L) & \text{if } i = 1, j \neq L \\ (1 - e(k, 0))/e(k, -L) & \text{if } i = 1, j = L \\ 1 & \text{if } i \geq 2, j = i - 1 \\ 0 & \text{otherwise} \end{cases}.$$

We also define the matrix $\mathbf{M} = \mathbf{A}_1 \mathbf{A}_2 \dots \mathbf{A}_n$.

According to (Key 1984) we define constants $d_i, i = 1, 2, \dots, R + L$ as follows. For each eigenvalue λ_i of $\mathbf{M}$ (including multiplicities), we put $d_i = \log(|\lambda_i|)$, and we list the d_i 's in increasing order, so that $d_1 \leq d_2 \leq \dots \leq d_{R+L}$. Then

- if $\ln(c(\mathbf{W})) \equiv d_R + d_{R+1} > 0$ then the RWPE X_n is transient to ∞ ;
- if $\ln(c(\mathbf{W})) \equiv d_R + d_{R+1} = 0$ then the RWPE X_n is recurrent ;
- if $\ln(c(\mathbf{W})) \equiv d_R + d_{R+1} < 0$ then the RWPE X_n is transient to $-\infty$. (2.2)

It is shown in (Key&Kłosek 2000) that X_n is recurrent if the characteristic polynomial of $\mathbf{M}$ has a double root at 1.

3. Games composed of 2-periodic games

First we consider an example of two 2-periodic games $\mathbf{P} = (P_0, P_1)$ and $\mathbf{Q} = (Q_0, Q_1)$; that is, we have two RWPE's, each of them 2-periodic (in space). If X_n denotes the capital of a player at time n which plays the game according to the rule $\mathbf{P}$, then the transition probabilities are given by

$$\begin{aligned} P\{X_{n+1} = x + 1 | X_n = x\} &= \begin{cases} p_0 & \text{if } x = 0 \bmod 2 \\ p_1 & \text{if } x = 1 \bmod 2 \end{cases} \\ P\{X_{n+1} = x - 1 | X_n = x\} &= \begin{cases} 1 - p_0 & \text{if } x = 0 \bmod 2 \\ 1 - p_1 & \text{if } x = 1 \bmod 2 \end{cases} ; \end{aligned} \quad (3.1)$$

that is, if the capital is even it changes according to P_0 , and it is odd it changes according to P_1 – cf. figure 1. The transition probabilities for the RWPE governed by $\mathbf{Q}$ are given by formulas analogous to (3.1), with p 's replaced by q 's, with the rule Q_0 and Q_1 at the even and odd positions, respectively.

To employ the criterion of (Key 1984) to the RWPE governed by $\mathbf{P}$ we construct the matrix $\mathbf{M}$

$$\mathbf{M} = \mathbf{A}_1 \mathbf{A}_0 \equiv \begin{bmatrix} 1/(1 - p_1) & -p_1/(1 - p_1) \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1/(1 - p_0) & -p_0/(1 - p_0) \\ 1 & 0 \end{bmatrix}.$$

Since here $R = L = 1$, $d_R + d_{R+1} = d_1 + d_2 = \log(|\det(\mathbf{M})|)$, so we find that the RWPE governed by $\mathbf{P}$ is transient to $-\infty$, recurrent, or transient to $+\infty$ if $c(\mathbf{P}) \equiv c(P_0, P_1) \equiv \frac{p_0 p_1}{(1 - p_0)(1 - p_1)} < (=)(>) 1$. We note that this analysis easily extends to any period N of the environment – cf. §4.




Figure 1. Schematic representation of a 2-(state) periodic game $\mathbf{P} = (P_0, P_1)$. Games composed of such games are discussed in §3.

(a) *An example of a pseudo-paradox*

We suppose that the games $\mathbf{P}$ and $\mathbf{Q}$ are fair, that is $c(P_0, P_1) = p_0(1 - p_0)/(p_1(1 - p_1)) = 1$ and $c(Q_0, Q_1) = q_0(1 - q_0)/(q_1(1 - q_1)) = 1$. If we alternate (deterministically) games $\mathbf{P}$ and $\mathbf{Q}$ in that very order, then, starting at the origin, the composite game may be fair (losing) (winning) if $c(\mathbf{P}, \mathbf{Q}) = p_0q_1/((1 - p_0)(1 - q_1)) = (<)(>)1$. Under this strategy we play the game P_0 when the winnings are even and the game Q_1 when the winnings are odd. The games P_1 and Q_0 are never played. Hence $c(\mathbf{P}, \mathbf{Q}) = c(P_0, Q_1)$, and the fact that the composite game is losing when it is constructed from two winning games is just an apparent paradox. If we play the same two games when starting at an odd position then $c(\mathbf{P}, \mathbf{Q}) = c(P_1, Q_0) = 1/c(P_0, Q_1)$, and we have a winning game constructed out of two winning games.

(b) *Effects of randomisation*

Now we construct a RWPE, X_n , by choosing at random between $\mathbf{P}$ and $\mathbf{Q}$. If our capital is even then a coin with probability of heads g_0 is used to choose whether the game P_0 or Q_0 will be played; if our capital is odd then a coin with probability of heads g_1 is used to choose between the games P_1 and Q_1 . In this scheme all four games are played. The transition probabilities are given by

$$P\{X_{n+1} = x + 1 | X_n = x\} = \begin{cases} g_0p_0 + (1 - g_0)q_0 & \text{if } x = 0 \bmod 2 \\ g_1p_1 + (1 - g_1)q_1 & \text{if } x = 1 \bmod 2 \end{cases} \quad (3.2)$$

The fairness of the composite game is determined by the factor $c(\mathbf{P}, \mathbf{Q})$ given by

$$c(\mathbf{P}, \mathbf{Q}) = \frac{g_0p_0 + (1 - g_0)q_0}{(1 - g_0p_0 - (1 - g_0)q_0)} \frac{g_1p_1 + (1 - g_1)q_1}{(1 - g_1p_1 - (1 - g_1)q_1)}. \quad (3.3)$$

By direct simplification of equation(3.3) we observe that if $g_0 = g_1$ (and the games $\mathbf{P}$ and $\mathbf{Q}$ are fair) then the composite game is fair. Also, if $p_1 = q_1$ then the composite game is fair. However, for other values of parameters two fair games can be used to compose a game which is winning, losing or fair depending on how the two simple games are randomised. Figure 2 illustrates this statement. That is, randomisation can produce Parrondo's paradox.




Figure 2. We take two 2-periodic fair games with $p_1 = 1/2$ and $q_1 = 1/4$, and plot $c(\mathbf{P}, \mathbf{Q}) - 1$ in Eq.(3.3) as a function of g_1 for various values of g_0 ; that is, the composite game is winning for positive values on the graph, and losing for negative values. We have: dashes: $g_0 = 0$; crosses: $g_0 = 1/8$; boxes: $g_0 = 1/2$; solid: $g_0 = 1$.

We also consider a game composed from two unfair games $\mathbf{P}$ and $\mathbf{Q}$. Games $\mathbf{P}$ and $\mathbf{Q}$ are taken to be unfair the same way; that is both are losing or both are winning. We note that when one coin is used to choose whether $\mathbf{P}$ or $\mathbf{Q}$ is played, (*i.e.*, when $g_0 = g_1$ in equation (3.2)) then the composed game is always unfair. However, if two coins are used then the randomised game may be winning when both simple games $\mathbf{P}$ and $\mathbf{Q}$ are losing. This example of Parrondo's paradox is illustrated in figure 3.

(c) Three 2-periodic games

We consider three 2-periodic games $\mathbf{P}$, $\mathbf{Q}$ and $\mathbf{R}$, each defined analogously to (3.1), with transition probabilities given in terms of p_0, p_1, q_0, q_1 , and r_0, r_1 , respectively. We construct the composite game $\mathbf{PQR}$. We investigate whether the composite game may be winning (losing) if the individual games $\mathbf{P}$, $\mathbf{Q}$ and $\mathbf{R}$ are fair. If the capital of the player at time n is X_n then the capital at times $3n$, $Y_n \equiv X_{3n}$, is a random walk in a 2-periodic environment, taking steps of $\pm 3, \pm 1$. Specifically, we have




Figure 3. We take two 2-periodic losing games with $p_0 = 0.675$, $p_1 = 0.1$, $q_1 = 0.75$ and plot the fairness curve $c(\mathbf{P}, \mathbf{Q}) = 1$ in Eq.(3.3) in the (g_1, g_0) plane for various values of q_0 : circles: $q_0 = 0$; dashes: $q_0 = 0.075$; boxes: $q_0 = 0.125$; solid: $q_0 = 0.175$; crosses: $q_0 = 0.225$; diamonds: $q_0 = 0.25$. The composite game is winning when g_0 and g_1 are selected above the fairness curve.

$$\begin{aligned}
 P\{Y_{n+1} = x + 3 | Y_n = x\} &= \begin{cases} a_0 \equiv p_0 q_1 r_0 & \text{if } x = 0 \bmod 2 \\ a_1 \equiv p_1 q_0 r_1 & \text{if } x = 1 \bmod 2 \end{cases} \\
 P\{Y_{n+1} = x - 3 | Y_n = x\} &= \begin{cases} b_0 \equiv (1 - p_0)(1 - q_1)(1 - r_0) & \text{if } x = 0 \bmod 2 \\ b_1 \equiv (1 - p_1)(1 - q_0)(1 - r_1) & \text{if } x = 1 \bmod 2 \end{cases} \\
 P\{Y_{n+1} = x + 1 | Y_n = x\} &= \\
 &\quad \begin{cases} c_0 \equiv p_0 q_1 (1 - r_0) + p_0 (1 - q_1) r_0 + (1 - p_0) q_1 r_0 & \text{if } x = 0 \bmod 2 \\ c_1 \equiv p_1 q_0 (1 - r_1) + p_1 (1 - q_0) r_1 + (1 - p_1) q_0 r_1 & \text{if } x = 1 \bmod 2 \end{cases} \\
 P\{Y_{n+1} = x - 1 | Y_n = x\} &= \\
 &\quad \begin{cases} d_0 \equiv (1 - p_0)(1 - q_1) r_0 + (1 - p_0) q_1 (1 - r_0) + p_0 (1 - q_1)(1 - r_0) & x = 0 \bmod 2 \\ d_1 \equiv (1 - p_1)(1 - q_0) r_1 + (1 - p_1) q_0 (1 - r_1) + p_1 (1 - q_0)(1 - r_1) & x = 1 \bmod 2. \end{cases}
 \end{aligned}$$

Moreover, the random walk $Z_n \equiv Y_{2n}$ is an ordinary random walk, that is a sum of iid random variables taking values $\pm 6, \pm 4, \pm 2$, and 0, and the transition probabilities are




Figure 4. Schematic representation of a 3-(state) periodic game $\mathbf{P} = (P_0, P_1, P_2)$. Games composed of such games are discussed in §4.

$$\begin{aligned}
 P\{Z_{n+1} = z + 6 | Z_n = z\} &= a_0 a_1 \\
 P\{Z_{n+1} = z - 6 | Z_n = z\} &= b_0 b_1 \\
 P\{Z_{n+1} = z + 4 | Z_n = z\} &= a_0 c_1 + c_0 a_1 \\
 P\{Z_{n+1} = z - 4 | Z_n = z\} &= b_0 d_1 + d_0 b_1 \\
 P\{Z_{n+1} = z + 2 | Z_n = z\} &= c_0 c_1 + a_0 d_1 + a_1 d_0 \\
 P\{Z_{n+1} = z - 2 | Z_n = z\} &= d_0 d_1 + b_0 c_1 + b_1 c_0 \\
 P\{Z_{n+1} = z | Z_n = z\} &= d_0 c_1 + d_1 c_0 + a_0 b_1 + a_1 b_0.
 \end{aligned}$$

If the games $\mathbf{P}$, $\mathbf{Q}$ and $\mathbf{R}$ are fair, then by direct calculations, $EZ_n = 0$, so the composite game is fair, and there is no paradox in this case.

4. Games composed of 3-periodic games

We consider games composed of 3-periodic games $\mathbf{P}$, $\mathbf{Q}$, and $\mathbf{R}$. We define a 3-periodic game $\mathbf{P} = (P_0, P_1, P_2)$ by its transition probabilities as

$$P\{X_{n+1} = x + 1 | X_n = x\} = \begin{cases} p_0 & \text{if } x = 0 \bmod 3 \\ p_1 & \text{if } x = 1 \bmod 3 \\ p_2 & \text{if } x = 2 \bmod 3 \end{cases}, \quad (4.1)$$

compare figure 4. Transition probabilities of the games $\mathbf{Q}$ and $\mathbf{R}$ are defined by formulas analogous to (4.1) with p 's replaced by q 's and r 's, respectively. According to (Key 1984), to determine the fairness of the game $\mathbf{P}$ we consider the product of the eigenvalues of the matrix $\mathbf{M} \equiv \mathbf{A}_1 \mathbf{A}_2 \mathbf{A}_0$ where

$$\mathbf{A}_i = \begin{bmatrix} 1/(1-p_i) & -p_i/(1-p_i) \\ 1 & 0 \end{bmatrix} \quad i = 0, 1, 2.$$

If $\frac{p_0 p_1 p_2}{(1-p_0)(1-p_1)(1-p_2)} = (<)(>)1$ then the game $\mathbf{P}$ is fair (losing) (winning).

(a) Two 3-periodic games

To analyse a game $(\mathbf{P}, \mathbf{Q})$, where $\mathbf{P}$ and $\mathbf{Q}$ are 3-periodic games, we construct a random walk $Y_n \equiv X_{2n}$ where X_n denotes the position of the walker at time n . The transition probabilities for Y_n are given by

$$P\{Y_{n+1} = x + 2 | Y_n = x\} = \begin{cases} a_0 \equiv p_0 q_1 & \text{if } x \equiv 0 \pmod{3} \\ a_1 \equiv p_1 q_2 & \text{if } x \equiv 1 \pmod{3} \\ a_2 \equiv p_2 q_0 & \text{if } x \equiv 2 \pmod{3} \end{cases}$$

$$P\{Y_{n+1} = x | Y_n = x\} = \begin{cases} b_0 \equiv p_0(1 - q_1) + (1 - p_0)q_2 & \text{if } x \equiv 0 \pmod{3} \\ b_1 \equiv p_1(1 - q_2) + (1 - p_1)q_0 & \text{if } x \equiv 1 \pmod{3} \\ b_2 \equiv p_2(1 - q_0) + (1 - p_2)q_1 & \text{if } x \equiv 2 \pmod{3} \end{cases}$$

$$P\{Y_{n+1} = x - 2 | Y_n = x\} = \begin{cases} c_0 \equiv (1 - p_0)(1 - q_2) & \text{if } x \equiv 0 \pmod{3} \\ c_1 \equiv (1 - p_1)(1 - q_0) & \text{if } x \equiv 1 \pmod{3} \\ c_2 \equiv (1 - p_2)(1 - q_1) & \text{if } x \equiv 2 \pmod{3} \end{cases}$$

Hence, Y_n is a 3-periodic random walk taking steps ± 2 and zero. To employ Key's criterion most efficiently we observe that it is sufficient to analyse Y_n as a RWPE on the even integers which visits only nearest neighbours. To this end we construct the 2×2 matrix $\mathbf{M} = \mathbf{A}_1 \mathbf{A}_2 \mathbf{A}_0$ where

$$\mathbf{A}_i = \begin{bmatrix} (1 - b_i)/c_i & -a_i/c_i \\ 1 & 0 \end{bmatrix} \quad i = 0, 1, 2.$$

The determinant of $\mathbf{M}$ is given by $\det(\mathbf{M}) = p_0 p_1 p_2 q_0 q_1 q_2 / [(1 - p_0)(1 - p_1)(1 - p_2)(1 - q_0)(1 - q_1)(1 - q_2)]$. Hence, if the games $\mathbf{P}$ and $\mathbf{Q}$ are fair, the composite game is fair and no paradox is observed.

We note that this reduction in the dimension of $\mathbf{M}$ occurs whenever the temporal period is even, since the step sizes of the derived process Y_n are even. Specifically, if the temporal period is T is even then $\mathbf{M}$ can be taken to be the product of $T \times T$ matrices, and when the temporal period T is odd, then $\mathbf{M}$ is the product of $2T \times 2T$ matrices.

(b) Two 3-periodic games randomised

We construct a composite game from two 3-periodic random games $\mathbf{P} = (P_0, P_1, P_2)$ and $\mathbf{Q} = (Q_0, Q_1, Q_2)$ by selecting at random the game to be played at each step, with probability of playing P_i equal to $g_i, i = 0, 1, 2$. That is, we define a 3-periodic random walk Y_n taking values ± 1 with transition probabilities

$$P\{Y_{n+1} = x + 1 | Y_n = x\} = \begin{cases} \rho_0 \equiv g_0 p_0 + (1 - g_0)q_0 & \text{if } x \equiv 0 \pmod{3} \\ \rho_1 \equiv g_1 p_1 + (1 - g_1)q_1 & \text{if } x \equiv 1 \pmod{3} \\ \rho_2 \equiv g_2 p_2 + (1 - g_2)q_2 & \text{if } x \equiv 2 \pmod{3} \end{cases} \quad (4.2)$$

The walk Y_n is recurrent (transient to ∞) (transient to $-\infty$) according to $c(\mathbf{P}, \mathbf{Q}) \equiv \frac{\rho_0 \rho_1 \rho_2}{(1 - \rho_0)(1 - \rho_1)(1 - \rho_2)} = (>)(<)1$. We investigate the value of $c(\mathbf{P}, \mathbf{Q})$




Figure 5. We take two fair 3-periodic games with $g_1 = g_2$, $p_1 = p_2 = 1/2$, and $q_1 = q_2$. For a fixed value of q_1 we plot curves along which the composite game is fair, in the space (g_1, g_0) . As q_1 varies from 0 to $1/2$ the composite game is losing above and winning below its corresponding “fairness” curve; solid: $q_1 = 0$; boxes: $q_1 = 1/6$; crosses: $q_1 = 1/3$; dashes: the line $g_0 = g_1$. As q_1 varies from $1/2$ to 1 the composite game is winning below and losing above its corresponding “fairness” curve. In this case the problem is symmetric, so that we have: solid: $q_1 = 1$; boxes $q_1 = 5/6$; crosses: $q_1 = 2/3$.

for various values of the parameters. We note that if the games $\mathbf{P}$ and $\mathbf{Q}$ are fair with $p_1 = p_2 = q_1 = q_2$ then the randomised composite game is fair for all possible choices of g_i 's. Next we take two fair games with fixed values $p_1 = p_2$ and $q_1 = q_2$, that is $\mathbf{P} = (P_0, P_1, P_1)$ and $\mathbf{Q} = (Q_0, Q_1, Q_1)$.

We investigate how randomisation affects the fairness of the composite game. As shown in figure 5 there is a wide range of parameters for which the randomised game is unfair, so that randomisation induces the paradox. We note that the range of the randomisation parameters g 's is affected by the way games $\mathbf{P}$ and $\mathbf{Q}$ are structured. In the examples illustrated in figure 6 we take two fair games $\mathbf{P} = (P_0, P_1, P_1)$ and $\mathbf{Q} = (Q_0, Q_1, Q_0)$ and we note that the range of parameters $g_0, g_1 = g_2$ for which the randomised game composed from these “shifted” games is larger compared to the previous “not-shifted” case – cf. figures 5 and 6.

(c) Three 3-periodic games

First we construct a game $(\mathbf{PQR})$ in which the 3-periodic games $\mathbf{P}$, $\mathbf{Q}$ and $\mathbf{R}$ are played in a deterministic order. We show that if all three games are fair then the resulting game may be losing. We also show that if all three games are winning then the composite game may be losing. If X_n denotes the winnings of the composite game at time n then its transition probabilities are given by




Figure 6. We take two fair 3-periodic games with $g_1 = g_2$, $p_1 = p_2 = 1/2$, and $q_0 = q_2$. We plot the curves along which the composite game is fair in the space (g_1, g_0) , for a fixed value of q_0 . As q_0 varies from 0 to $1/2$, the game is winning above and losing below each of its “fairness” curves; solid: $q_0 = 0$; circles: $q_0 = 1/12$; boxes: $q_0 = 1/6$; crosses: $q_0 = 1/3$; dashes: the line $g_0 = g_1$. As q_0 increases from $1/2$ to 1 the game is losing above and winning below its fairness curve. The case of this example is symmetric so that we have: crosses: $q_0 = 2/3$; boxes: $q_0 = 5/6$; circles: $q_0 = 11/12$.

if $x = 0 \bmod 3$ then

$$\begin{aligned}
 P\{X_{n+3} = x + 3 | X_n = x\} &= a_0 \equiv p_0 q_1 r_2 \\
 P\{X_{n+3} = x - 3 | X_n = x\} &= b_0 \equiv (1 - p_0)(1 - q_2)(1 - r_1) \\
 P\{X_{n+3} = x + 1 | X_n = x\} &= c_0 \equiv \\
 &\quad p_0(1 - q_2)r_1 + (1 - p_0)q_2(1 - r_0) + p_0(1 - q_1)(1 - r_0) \\
 P\{X_{n+3} = x - 1 | X_n = x\} &= d_0 \equiv \\
 &\quad (1 - p_0)(1 - q_2)r_1 + (1 - p_0)q_2(1 - r_0) + p_0(1 - q_1)(1 - r_0)
 \end{aligned}$$

if $x = 1 \bmod 3$ then

$$\begin{aligned}
 P\{X_{n+3} = x + 3 | X_n = x\} &= a_1 \equiv p_1 q_2 r_0 \\
 P\{X_{n+3} = x - 3 | X_n = x\} &= b_1 \equiv (1 - p_1)(1 - q_0)(1 - r_2) \\
 P\{X_{n+3} = x + 1 | X_n = x\} &= c_1 \equiv p_1 q_2(1 - r_0) + p_1(1 - q_2)r_1 + (1 - p_1)q_0 r_1 \\
 P\{X_{n+3} = x - 1 | X_n = x\} &= d_1 \equiv \\
 &\quad (1 - p_1)(1 - q_0)r_2 + (1 - p_1)q_0(1 - r_1) + p_1(1 - q_2)(1 - r_1)
 \end{aligned}$$




Figure 7. We take 3 fair games with $p_0 = p_1 \equiv p$, $q_0 = q_1 \equiv q$, and $r_0 = r_1 \equiv r$, and plot $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R})) = d_3 + d_4$ – compare Eq.(2.2), as a function of p , for $q = 0.1$ and various values of r , solid: $r = 3/4$; crosses: $r = 1/2$, boxes: $r = 1/4$; short dashes: $r = 1/8$; long dashes: $r = 1/16$. We note that composite game is always losing, no matter what p is, for a wide range of values of the parameter r . If $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R}))$ is positive (zero) (negative) then the game $(\mathbf{P}, \mathbf{Q}, \mathbf{R})$ is winning (fair) (losing).




Figure 8. We take 3 fair games with $p_0 = p_1 \equiv p$, $q_0 = q_1 \equiv q$, and $r_0 = r_1 \equiv r$, and plot $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R})) = d_3 + d_4$ – compare (2.2), as a function of p , for $q = 3/4$ and various values of r : circles: $r = 3/4$, crosses: $r = 5/8$; dashes: $r = 1/2$; solid: $r = 3/8$. If $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R}))$ is positive (zero) (negative) then the game $(\mathbf{P}, \mathbf{Q}, \mathbf{R})$ is winning (fair) (losing).




Figure 9. We take 3 losing games with $p_0 = p_1 \equiv p$, $q_0 = q_1 \equiv q$, and $r_0 = r_1 \equiv r$, and $p_2 = 0.8(1 - p)^2 / ((1 - p)^2 + p^2)$, $q_2 = 0.8(1 - q)^2 / ((1 - q)^2 + q^2)$, and $r_2 = 0.9(1 - r)^2 / ((1 - r)^2 + r^2)$. We set $q = 3/4$, so that $q_2 = 0.08$, and plot $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R})) = d_3 + d_4$ – compare (2.2), as a function of p for various values of r , circles: $r = 1/2$, ($r_2 = 0.45$); solid: $r = 5/8$, ($r_2 = 0.238235 \dots$); dashes: $r = 3/4$, ($r_2 = 0.09$); crosses $r = 7/8$, ($r_2 = 0.018$). If $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R}))$ is positive (zero) (negative) then the game $(\mathbf{P}, \mathbf{Q}, \mathbf{R})$ is winning (fair) (losing).

if $x = 2 \bmod 3$ then

$$P\{X_{n+3} = x + 3 | X_n = x\} = p_2 q_0 r_1 \equiv a_2$$

$$P\{X_{n+3} = x - 3 | X_n = x\} = (1 - p_2)(1 - q_1)(1 - r_0) \equiv b_2$$

$$P\{X_{n+3} = x + 1 | X_n = x\} = p_2 q_0 (1 - r_1) + p_2 (1 - q_0) r_2 + (1 - p_2) q_1 r_2 \equiv c_2$$

$$P\{X_{n+3} = x - 1 | X_n = x\} = d_2 \equiv$$

$$(1 - p_2)(1 - q_1) r_0 + (1 - p_2) q_1 (1 - r_2) + p_2 (1 - q_0) (1 - r_2).$$

Hence we define the walk in periodic environment by $Y_n \equiv X_{3n}$, and determine its transient/recurrent properties by investigating locations of the eigenvalues of the matrix $\mathbf{M} = \mathbf{A}_1 \mathbf{A}_2 \mathbf{A}_0$. Since the walk Y_n takes on the values $\pm 3, \pm 1$, $R = L = 3$, and each of the matrices $\mathbf{A}_i, i = 1, 2, 3$ is a 6×6 matrix given by

$$\mathbf{A}_i = \begin{bmatrix} 0 & -d_i/b_i & 1/b_i & -c_i/b_i & 0 & -a_i/b_i \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}, i = 0, 1, 2.$$




Figure 10. We take 3 losing games with $p_0 = p_1 \equiv p$, $q_0 = q_1 \equiv q$, and $r_0 = r_1 \equiv r$, and $p_2 = 0.8(1-p)^2/((1-p)^2 + p^2)$, $q_2 = 0.8(1-q)^2/((1-q)^2 + q^2)$, and $r_2 = 0.9(1-r)^2/((1-r)^2 + r^2)$. We set $q = 0.1$ so that $q_2 = 0.8890\dots$, and plot $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R})) = d_3 + d_4 - \text{compare (2.2)}$, as a function of p for various values of r , solid: $r = 3/4$, ($r_2 = 0.9$); box: $r = 1/2$, ($r_2 = 0.45$); circles: $r = 3/8$, ($r_2 = 0.66176\dots$); crosses $r = 1/4$, ($r_2 = 0.81$). If $\ln(c(\mathbf{P}, \mathbf{Q}, \mathbf{R}))$ is positive (zero) (negative) then the game $(\mathbf{P}, \mathbf{Q}, \mathbf{R})$ is winning (fair) (losing).

The examples illustrated in figures 7 and 8 show that the game composed deterministically from fair (non-symmetric) games can be fair or losing or winning, depending on parameters of the problem. This form of the paradox is not observed when composing (deterministically) two or three 2-periodic games. We observe even more interesting behaviour when composing deterministically unfair (losing) games. Figures 9 and 10 show such examples.

By varying just one parameter, it is possible to change the composed game from losing to winning, change it back to losing, and then back again to winning – cf. figure 9. Three state transitions of the composed game have not been observed before.

5. Conclusions

Given the highly non-linear nature of the recurrence/transience criteria for random walks in periodic environments as a function of the process parameters, it is not surprising that any scheme that combines recurrent random walks in periodic environment to produce new random walks in periodic environments will produce transient processes, except in trivial cases.

What is surprising to us is that one can find non-trivial cases where random walks in periodic environments with negative drift can be combined to form random walks in periodic environments with positive drift, Parrondo's paradox. We believe that these reversals arise because combining these processes in the manner

described above radically changes the frequency at which transitions are governed by probability distributions with positive means, and that future investigations should center on investigating these frequencies. It is likely that it will be easier to understand this phenomenon in the case of stochastic combinations of the processes since these always lead to the study of nearest neighbour random walks in periodic environments.

When a composite game is constructed from simple 3-periodic games, Parrondo's paradox is much more interesting and complex than in the case of 2-periodic games. As shown in figure 9, by varying one parameter three state transitions of the composed game may be observed. This phenomenon seems to be similar to changes of the direction of the current in a multiplicative stochastic ratchet as observed in (Kłosek & Cox 2000).

As shown in §3, a game constructed by composing deterministically 2-periodic fair games leads to a pseudo-paradox. In general, a game composed from an even number of 2-periodic fair games always exhibits a pseudo-paradox (Key & Kłosek 2000). That is, if each simple game is 2-state periodic with an even temporal period, then the composite game is constructed only from half of the simple games, and the composite game may be winning, or losing, or fair. Moreover, when each of the simple games is fair with an odd temporal period, and the composite game is composed deterministically, then Parrondo's paradox never occurs. An example in §3 of a game composed from 2-state periodic and 3-temporal periodic games illustrates this statement.

D.A. would like to acknowledge funding from the Sir Ross and Sir Keith Smith Fund, GTECH, the Australian Research Council and SA Lotteries. Also thanks are due to Prof. A.E. Seigman, Stanford University, USA and Prof. J. Maynard Smith, University of Sussex, UK for useful communications. The 'Boston Interpretation' originated during a round table discussion with Prof. H.E. Stanley's group at Boston University.

References

- Allison, A. & Abbott, D. 2000 Stochastically switched control systems. *Proc. 2nd Int. Conf. on Unsolved Problems of Noise (UPoN'99)*, Adelaide, Australia, 11-15th July 1999, pp. xx-xx. American Institute of Physics (In Press).
- Doering, C.R. 1995 Randomly rattled ratchets. *Nuovo Cimento*. **17D** 685-679.
- Eisert, J., Wilkens, M., and Lewenstein, M. 1999 Quantum games and quantum strategies *Phys. Rev. Lett.*, **83**, 3077-3080.
- Harmer, G.P. and Abbott, D. 1999a Losing strategies can win by Parrondo's paradox. *Nature*. **402**, 846.
- Harmer, G.P. and Abbott, D. 1999b Parrondo's paradox. *Statistical Science*. **14**, 2, 206-213.
- Harmer, G.P., Abbott, D., Taylor, P. & Parrondo, J.M.R. 2000a Parrondo's paradoxical games and the discrete Brownian ratchet. *Proc. 2nd Int. Conf. on Unsolved Problems of Noise (UPoN'99)*, Adelaide, Australia, 11-15th July 1999, pp. xx-xx. American Institute of Physics (In Press).
- Harmer, G.P., Abbott, D., Taylor, P., Pearce, C.E.M. & Parrondo, J.M.R. 2000b Information entropy and Parrondo's discrete-time ratchet, *Proc. Stochaos, Ambleside, UK, 16-20 August 1999*, pp. xx-xx. American Institute of Physics (In Press).
- Harmer, G.P., Abbott, D. & Taylor, P. 2000c The paradox of Parrondo's games. *Proc. R. Soc. Lond. A*. xx-xx. (In Press).

- Gargamo, L. & Vaccaro, U. 1999 Efficient generation of fair dice with few biased coins. *IEEE Trans. Info. Theory*, **45**, 5, 1600-1606.
- Goldenberg, L., Vaidman, L., and Wiesner, S. 1999 Quantum gambling *Phys. Rev. Lett.*, **82** 3356-3359.
- Key, E.S. 1987 Computable Examples of the Maximal Lyapunov Exponent. *Probab. Th. Rel. Fields* **75**, 97-107.
- Key, E.S. 1984 Recurrence and Transience Criteria for Random Walk in a Random Environment. *Ann. Prob.* **12**, 529-560.
- Key, E.S. & Kłosek, M.M. 2000 Temporally periodic random walks in periodic environments. (In preparation).
- Kłosek, M.M. & Cox, R.W. 2000 Steady-State Currents in Sharp Stochastic Ratchets, *Proc. Stochaos, Ambleside, UK, 16-20 August 1999*, pp. xx-xx. American Institute of Physics (In Press).
- Lee, Y., Abbott, D. & Stanley, H.E. 2000 Minimal Parrondian ratchet: an exactly solvable model. *Phys. Rev. Lett.* xx-xx. (Submitted).
- Linke, H., Humphrey, T.E., Löfgren, A., Sushkov, A.O., Newbury, R., Taylor, R.P., and Omling, P. 1999 Experimental tunneling ratchets *Science*, **286** 2314-2317.
- Maslov, S. & Zhang, Y. 1998 Optimal investment strategy for risky assets. *Int. J. of Theoretical and Appl. Finance*, **1**, 377-387.
- McClintock, P.V.E. 1999 Unsolved problems of noise. *Nature*, **401** 23-25.
- Meyer, D.A. 1998 Quantum strategies *Phys. Rev. Lett.*, **82** 1052-1055.
- Parrondo, J.M.R., Harmer, G.P. & Abbott, D. 2000 New paradoxical games based on Brownian ratchets. *Phys. Rev. Lett.* xx-xx. (Submitted).
- Pearce, C.E.M. 2000a On Parrondo's paradoxical games. *Proc. 2nd Int. Conf. on Unsolved Problems of Noise (UPoN'99), Adelaide, Australia, 11-15th July 1999*, pp. xx-xx. American Institute of Physics (In Press).
- Pearce, C.E.M. 2000b Entropy, Markov information sources and Parrondo games. *Proc. 2nd Int. Conf. on Unsolved Problems of Noise (UPoN'99), Adelaide, Australia, 11-15th July 1999*, pp. xx-xx. American Institute of Physics (In Press).
- Pinsky, R., and Scheutzow, M. 1992 Some remarks and examples concerning transience and recurrence of random diffusions, *Ann. Inst. H. Poincaré Probab. Statist.*, **28** 519-536.
- Plaskota, L. 1996 How to benefit from noise. *J. Complexity*, **12**, 175-184.
- Rosato, A., Strandburg, K.J., Prinz, F., Swendsen, R.H. 1987 Why the Brazil nuts are on top: size segregation of particulate matter shaking. *Phys. Rev. Lett.*, **58** 1038-1040.
- Sarmiento, A., Reigada, R., Romero, A.H., & Lindenberg, K. 1999 Enhanced pulse propagation in non-linear arrays of oscillators. *Phys. Rev. E*, **60** 5317-5326
- von Neumann, J. 1951 Various techniques used in connection with random digits, notes by G.E. Forsythe, National Bureau of Standards. *Appl. Math. Ser.* **12**, 36-38.
- Westerhoff, H.V., Tsong, T.Y., Chock, P.B., Chen, Y. and Astumian, R.D. 1986 How enzymes can capture and transmit free energy from an oscillating electric field. *Proc. Natl. Acad. Sci.*, **83** 4734-4738.